

## WEST

# Freeform Search

Database:	US Patents Full-Text Database  JPO Abstracts Database  EPO Abstracts Database  Derwent World Patents Index  IBM Technical Disclosure Bulletins			
Term:	16 same 12			
Display: Generate:	Documents in <u>Display Format</u> : TI Starting with Number 1 O Hit List O Hit Count O Image			
-				
	Search Clear Help Logout Interrupt			
	Main Menu Show S Numbers Edit S Numbers Preferences			

### Search History

Today's Date: 8/8/2000

<u>DB Name</u>	Query	Hit Count	Set Name
USPT,JPAB,EPAB,DWPI,TDBD	16 same 12	3	<u>L7</u>
USPT,JPAB,EPAB,DWPI,TDBD	rate near2 (reaction or reactions)	53462	<u>L6</u>
USPT,JPAB,EPAB,DWPI,TDBD	12 same 14	4	<u>L5</u>
USPT,JPAB,EPAB,DWPI,TDBD	rate near5 (measurement or measurements)	22822	<u>L4</u>
USPT,JPAB,EPAB,DWPI,TDBD	11 and 12	28	<u>L3</u>
USPT,JPAB,EPAB,DWPI,TDBD	electrochemiluminescence or electrochemiluminescent	389	<u>L2</u>
USPT,JPAB,EPAB,DWPI,TDBD	(kinetic or kinetics) same (reaction or reactions)	12111	<u>L1</u>

### End of Result Set

Generate Collection

L5: Entry 4 of 4

File: EPAB

Aug 26, 1992

PUB-NO: EP000500305A2

DOCUMENT-IDENTIFIER: EP 500305 A2

TITLE: Method of measuring immunoreactant using electro-chemiluminescence.

PUBN-DATE: August 26, 1992

INVENTOR-INFORMATION:

NAME COUNTRY
SHIBUE, AKIRA JP
TANAKA, MASARU JP
KAMIYA, SHINJI JP

ASSIGNEE-INFORMATION:

NAME COUNTRY

TDK CORP

APPL-NO: EP92301295

APPL-DATE: February 18, 1992

PRIORITY-DATA: JP04605491A (February 19, 1991)

US-CL-CURRENT: 436/172; 436/536

INT-CL (IPC): GOIN 21/66; GOIN 33/533; GOIN 33/546; GOIN 33/68

EUR-CL (EPC): G01N033/543; G01N033/58, G01N033/68

#### ABSTRACT:

A method of measuring an immnoreactant in a liquid sample, comprising the steps of: mixing the liquid sample with an excess of a complementary immunoreactant capable of specifically binding to said immunoreactant to allow an immunoreaction to take place, said complementary immunoreactant having been immobilized on insoluble carrier particles and labeled with an electrochemiluminescent substance that emit an electrochemiluminescent light by electrolytic oxidation in the presence of an activated oxygen, applying an electric voltage to a pair of electrodes between which the mixture obtained above is placed, in the presence of an activated oxygen to allow electrochemiluminescence to take place, and measuring the emission of the electrochemiluminescent light. The accuracy of measurement is high since the rate of change of emission of luminescent light depending on the

JP

change of concentration of the immunoreactant in the liquid sample is high.

8/8/00 2:07 PM